

## PACKAGE LEAFLET:

### INFORMATION FOR THE USER

**Dioralyte Natural/Citrus/Blackcurrant**

**3.5g/470mg/300mg/530mg,**

**Powder for oral solution**

Glucose, sodium chloride, potassium chloride,  
disodium hydrogen citrate

**SANOFI** 

**Is this leaflet hard to see or read?**

**Phone 01 403 5600 for help**

 **Read all of this leaflet carefully because it contains important information for you.**

This medicine is available without prescription. However, you still need to use Dioralyte Sachets carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must contact a doctor if your symptoms worsen or do not improve after using this medicine for 24-48 hours.
- If you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

#### **In this leaflet:**

1. What Dioralyte is and what it is used for
2. Before you take Dioralyte
3. How to take Dioralyte
4. Possible side effects
5. How to store Dioralyte
6. Further information

## **1. What Dioralyte is and what it is used for**



The name of your medicine is Dioralyte Natural/Citrus/Blackcurrant (called Dioralyte throughout this leaflet). Dioralyte works by replacing the water and salts lost from your body when you have diarrhoea. It helps the watery stools return to normal.

It contains glucose and salts (these are sodium chloride, potassium chloride and disodium hydrogen citrate). The powder is mixed with water before taking.

## **2. Before you take Dioralyte**

 **Do not take Dioralyte and talk to your doctor or pharmacist if:**

- × You are allergic (hypersensitive) to glucose, sodium chloride, potassium chloride and disodium hydrogen citrate or any of the other ingredients of Dioralyte (see Section 6: Further information). Signs of an allergic reaction include: a rash, swallowing or breathing problems, swelling of your lips, face, throat or tongue
- × Your bowel is partially or fully blocked (called 'intestinal obstruction')

Do not take this medicine if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Dioralyte.



**Take special care with Dioralyte. Check with your doctor or pharmacist before taking this medicine if:**

- ▲ The child taking the medicine is under 2 years old
- ▲ The diarrhoea has lasted for more than 24 to 36 hours, you may need other treatment
- ▲ You are on a low potassium or low sodium diet
- ▲ You have liver or kidney disease
- ▲ You have diabetes

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Dioralyte Relief.



#### **Taking other medicines**

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines. This includes medicines you buy without a prescription, including herbal medicines. This is because Dioralyte can affect the way some other medicines work. Also some medicines can affect the way Dioralyte works.

#### **Pregnancy and breast-feeding**

Ask your doctor or pharmacist for advice before taking any medicine if you are pregnant or breast-feeding. This includes if you are planning to get pregnant or think you that you might be pregnant.

#### **Important information about some of the ingredients of Dioralyte**

This medicine contains glucose, which is a type of sugar. If you have been told by your doctor that you cannot tolerate or digest some sugars, talk to your doctor before taking this medicine

## **3. How to take Dioralyte**

If you are not sure how to take this medicine, check with your doctor or pharmacist




#### **Preparing this medicine**

Dioralyte must be mixed with water before taking.

- Prepare just before the medicine is going to be taken. Do not make in advance. If you are giving it to a baby over the course of the day, you can keep the drink in a refrigerator for up to 24 hours
- **Only** dissolve the powder in water. Do not dissolve it in anything else
- Use fresh drinking water for adults and children over 1 year old. If fresh water is not available, use water which has been freshly boiled and cooled
- Always use water which has been freshly boiled and cooled for babies aged 3 months to 1 year







1 Empty the contents of 1 sachet into a glass

2 Add 200 ml (7 fl oz) of water

3 Stir until the powder dissolves

When your medicine is fully dissolved, the drink will be a colourless solution.

Do not use more or less than 200 ml of water. If you do, the drink will become too weak or too strong. It will not work as well. You may also upset the balance of salts in your body.

#### How many sachets to take

- If you are being sick take the medicine in small sips
- It is best not to eat during the early stages of diarrhoea
- If the diarrhoea lasts for more than 24 to 36 hours see your doctor

#### Adults (including the elderly)

- One or two sachets after each loose stool

#### Children

- One sachet after each loose stool

#### Infants ages 1 - 2 years

**Infants under 1 year: Use only on medical advice**

**This medicine should only be given under the guidance of a doctor**

- The doctor will calculate how much to give your baby, depending on how much they weigh. This is usually about one to one and a half times the amount of fluid your baby normally has in 24 hours
- Initially, if you are bottle feeding or giving other foods this should be stopped
- If you are breast feeding, continue to feed normally. Your baby may need less breast milk so you may need to express some of your milk
- Ask your doctor or pharmacist if you are not sure how much to give
- **If your baby is being sick**, you can give 5-10 ml (one or two teaspoons) every 5 minutes. This can be gradually increased until they are able to drink normally

#### If you take more Dioralyte than you should

Dioralyte is not normally harmful if you take more than you should. However, if large amounts are taken, you should talk to a doctor or go to a hospital straight away. This is particularly important if it is a child or baby or if you have severe liver or kidney problems. Take the medicine pack with you.

#### If you forget to take your Dioralyte

Do not take a double dose to make up for a forgotten dose. Take the next dose when you need to.



If you have any further questions on the use of this product, ask your doctor or pharmacist.

#### 4. Possible side effects

Like all medicines, Dioralyte can cause side effects, although not everybody gets them. However, if you notice anything unusual, talk to your doctor or pharmacist. You should not normally have any side effects.

**Stop taking Dioralyte and see a doctor or go to a hospital straight away if:**

- You have an allergic reaction. The signs may include: a rash, swallowing or breathing problems, swelling of your lips, face, throat or tongue

#### 5. How to store Dioralyte

Keep out of the reach and sight of children.

Do not use Dioralyte after the expiry date which is stated on the carton and sachet after EXP. The expiry date refers to the last day of that month.

Store below 25°C. Store in the original package in order to protect from moisture.

The drink should be made up just before it is going to be taken. Throw away any remaining drink after 1 hour (or 24 hours if stored in a refrigerator).

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

#### 6. Further information

##### What Dioralyte contains

- The active substances are glucose, sodium chloride, potassium chloride and disodium hydrogen citrate
- other ingredients include: colloidal anhydrous silica, saccharin sodium, lemon/lime flavour (contains sulphur dioxide (E220), citrus packs only), blackcurrant flavour (contains ethanol, blackcurrant packs only).  
Dioralyte Blackcurrant Powder contains small amounts of ethanol (alcohol), less than 100mg per dose.  
Dioralyte Citrus Powder contains sulphur dioxide. Sulphur dioxide may rarely cause severe hypersensitivity reactions and bronchospasm.

##### What Dioralyte looks like and contents of the pack

Dioralyte® Natural/Citrus/Blackcurrant is supplied in foil/laminate sachets available in pack sizes of 6 and 20. Each sachet contains enough powder to make a drink up with 200 ml (7 fl oz) of water.

##### Marketing Authorisation Holder and Manufacturer

###### Marketing Authorisation Holder

Sanofi-aventis Ireland Limited T/A SANOFI  
Citywest Business Campus, Dublin 24, Ireland.  
Tel: 01 403 5600, Fax: 01 403 5601  
email: IEmedinfo@sanofi-aventis.com

###### Manufacturer

A Nattermann & Cie. GmbH, Nattermannallee 1  
D-50829 Cologne, Germany.

This leaflet does not contain all the information about your medicine. If you have any questions or are not sure about anything, ask your doctor or pharmacist.

**This leaflet was last revised in {06/2012}.**

© Sanofi-aventis, 1995-2012

416770-05.12-IRL