FRONT LEADING EDGE



Film-coated tablets

Please read right through this leaflet before

This medicine is available without prescription,

but you still need to use Panadol tablets

carefully to get the best results from them.

Keep this leaflet, you may need to read

If you have any further questions, ask your

1. What Panadol tablets are and what they are

2. Check before you take Panadol tablets

5. How to store and dispose of Panadol

3. How to take Panadol tablets 4. Possible side effects

500mg

pharmacist.

In this leaflet:

used for

tablets

6. Further information

they are used for

Paracetamol

PATIENT INFORMATION LEAFLET

you start using this medicine.

2. Check before you take Panadol



Do not take Panadol tablets:

- if you have ever had an allergic reaction to paracetamol or to any of the other ingredients (listed in Section 6).
- if you are taking other medicines containing paracetamol.
- if you are under 6 years unless your doctor tells you to.



Ask your doctor before you take this

if you have severe liver or kidney disease, including alcoholic liver disease.



If you are taking other medicines

Talk to your doctor or pharmacist before taking these tablets if you are taking any prescribed medicines; particularly metoclopramide or domperidone (for nausea [feeling sick] or vomiting [being sick]); colestyramine (to lower **blood cholesterol**). If you take **blood** thinning drugs (anticoagulants e.g. warfarin) and you need to take a pain reliever on a daily basis, talk to your doctor because of the risk of bleeding. However you can still take occasional doses of Panadol tablets at the same time as anticoagulants.



Pregnancy and breastfeeding

Talk to your doctor before taking Panadol tablets if you are **pregnant.** You can take this product whilst breastfeeding.



Important information about some of the ingredients

Contains parahydroxybenzoates which may cause allergic reaction (possibly delayed).

3. How to take Panadol tablets



Adults, including the elderly, and children aged 12 years and over:

Swallow 2 tablets with water 3 to 4 times a day as needed. Do not take more frequently than every 4 hours. Do not take more than 8 tablets in 24 hours.

DEVCOMP-0003660

Panadol tablets are used for fast and

1. What Panadol tablets are and what

effective relief of headaches, toothache, backache, rheumatic and muscle pains and period pain. It also relieves sore throat and the feverishness, aches and pains of colds and flu. Panadol tablets also help relieve the pain of osteoarthritis (which has been diagnosed by a

The active ingredient is paracetamol which is a painkiller and also reduces your temperature when you have a fever.

TRAILING EDGE

TEXT SIZE CONTAINED IN THIS ARTWORK Body text size: 7.5pt Leading: 9.0pt **Horizontal Scale: 100%** Smallest text size: 7.5pt Microtext: N

	Dungarvan - Additional Artwork Information Panel	
	Pharmacode No.	N/A
7	Varnish Type	N/A
	Generic Specification Reference No.	N/A

GlaxoSmithKline **Artwork Information Panel**

RSC A/W

, THEN this document has NOT been printed from the Global Pack Manage

180 mm Measuring Bar

Item Number: **DEVCOMP-0003660**

Manufacturing Site:

GSK-IRL-Dungarvan (OTC)-IEDUN

Market or Pack Owner:

CH-Ireland-IRL; CH-UK Regulatory-GBR

Market Trade Name: Panadol Advance

Colour Standard Reference Number: N/A

Technical Reference No(s).: OTC.LT.002

do NOT include the technical reference doc[s] version no[s].)

Printing Process: Information Not Available

Substrate:

Κ

Information Not Available Colours Total: 1

Varnishes Total: N/A

Special Finishes Total: N/A

Artwork copyright is the property of the GlaxoSmithKline Group of Companies

All suppliers providing a service to GSK for printed components of any description must ensure that they have a licence for all fonts / software used in conjunction with GSK artwork.

The distribution and use of fonts / software without

a licence constitutes an intellectual property infringement. GSK will not accept any liability for the breach of third party intellectual property rights by printed component suppliers.

The GSK certification / audit process requires supplier to declare that they do not use unlicensed fonts / coffuses and may require the venture for the confuser.

software and may require the supplier to produce evidence of such licence to GSK.

ATTENTION • ATTENTION

To Ensure Accurate PDF Viewing and Printing: FOR SCREEN VIEWING: Use Adobe Acrobat 7 Professional or Adobe Acrobat Reader, Standard or Professional (higher than 7). **Overprint Preview** must be activated for accurate on screen viewing. FOR PRINTING: Use only Acrobat Professional rersion 7 or higher. "Apply Overprint Preview" or "Simulate Overprinting" must be activated in the print settings for printing accurate hard copies.

GSK Market is responsible for this product, its design and content. **Ensure the artwork is** thoroughly checked, all the text proof-read and approved.

RSC GSK is responsible for site technical requirements and pre-press suitability.

is responsible to advise RSC in case changes required impact the followings:

Formulation Tablet embossing Storage conditions **Shelf Life**

Dungarvan Additional Information Panel Reel Unwind Direction N/A

RSC A/W

CH-Ireland-IRL; CH-UK Regulatory-GBR

Panadol Advance

Colour Standard Reference

Technical Reference No(s).: OTC.LT.002 do NOT include the technical reference doc[s] version no[s].)

Printing Process:

Information Not Available

Substrate: **Information Not Available**

Colours Total: 1 Κ **Varnishes** Total: N/A

Special Finishes

Artwork copyright is the property of the GlaxoSmithKline Group of Companies

ATTENTION • ATTENTION

To Ensure Accurate PDF Viewing and Printing: FOR SCREEN VIEWING: Use Adobe Acrobat 7 Professional or Adobe Acrobat Reader, Standard or Professional (higher than 7). **Overprint Preview** must be activated for accurate on screen viewing. FOR PRINTING: Use only Acrobat Professional

GSK Market is responsible for this product, its design and content. **Ensure the artwork is** thoroughly checked, all the

GlaxoSmithKline **Artwork Information Panel**

Item Number: **DEVCOMP-0003660**

Manufacturing Site: GSK-IRL-Dungarvan (OTC)-IEDUN

Market or Pack Owner:

Market Trade Name:

Number: N/A

Total: N/A

All suppliers providing a service to GSK for printed components of any description must ensure that they have a licence for all fonts / software used in conjunction with GSK artwork.

The distribution and use of fonts / software without

a licence constitutes an intellectual property infringement. GSK will not accept any liability for the breach of third party intellectual property rights by printed component suppliers.

The GSK certification / audit process requires suppliers to declare that they do not use unlicensed fonts / coffuses and may require the certification for the confuse of the confu

software and may require the supplier to produce evidence of such licence to GSK.

version 7 or higher. "Apply Overprint Preview" or "Simulate Overprinting" must be activated in the print settings for printing accurate hard copies.

LEADING EDGE

BACK

Children aged 6-12 years: Give 1/2 to 1 tablet to swallow 3-4 times daily as required. Do not give more frequently than every 4 hours.

Do not give more than 4 tablets in 24 hours.

Not recommended for children under

6 years except on medical advice



- Do not take for more than three days without consulting your doctor.
- · Do not take more than the recommended dose.
- If symptoms persist consult your doctor.
- Prolonged use except under medical supervision may be harmful.

If you take too many tablets

Immediate medical advice should be sought in the event of an overdose, even if you feel well, because of the risk of delayed, serious

4. Possible side effects

Like all medicines Panadol tablets can have side effects, but not everybody gets them:

Stop taking this medicine and tell your doctor immediately if:

- you experience allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face.
- you experience a severe skin rash or peeling of the skin which may be accompanied by mouth ulcers.
- you may have previously experienced breathing problems with aspirin or non-steroidal anti-inflammatories, and experience a similar reaction to this product.
- you experience unexplained bruising or bleeding.

These reactions are rare.

Reporting of side effects If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store and dispose of **Panadol tablets**

Keep out of the sight and reach of children. This medicinal product does not require any special storage conditions.

Do not use this medicine after the expiry date (EXP) which is stated on the outer carton and the blisters. The expiry date refers to the last day of that month.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Further information

Panadol tablets contains the Optizorb™ formulation:

Active ingredient Each tablet contains Paracetamol 500mg.

Other ingredients Pregelatinised starch, calcium carbonate, alginic acid, crospovidone, povidone (K-25), magnesium stearate, colloidal anhydrous silica, parahydroxybenzoates (sodium methyl parahydroxybenzoate [E219], sodium ethyl parahydroxybenzoate [E215] and sodium propyl parahydroxybenzoate [E217]), opadry white (YS-1-7003), which contains titanium dioxide, hypromellose, macrogol, polysorbate 80 and carnauba wax.

Panadol tablets are white, capsule shaped, filmcoated tablets with a "P" within a circle on one face and a breakline on the other. The tablet can be divided into two equal halves.

Panadol tablets are produced in packs of 6, 12, 24, 48 or 96 tablets. Not all pack sizes may be

The Product Authorisation holder is GlaxoSmithKline Consumer Healthcare (Ireland) Limited, Stonemasons Way, Rathfarnham, Dublin 16 and all enquiries

should be sent to this address. The manufacturer is GlaxoSmithKline Dungarvan Ltd., Knockbrack, Dungarvan,

Co. Waterford, Ireland.

This leaflet was last revised in August 2014. **Panadol** is a registered trade mark of the GSK

PA 678/107/1

group of companies.

DEVCOMP-0003660

TRAILING EDGE

TEXT SIZE CONTAINED IN THIS ARTWORK		
Body text size: 7.5pt		
Leading: 9.0pt		
Horizontal Scale: 100%		
Smallest text size: 7.5pt		
Microtext: N		

Pharma code Ref. No. XXX

Dungarvan - Additional Artwork Information Panel	
Pharmacode No.	N/A
Varnish Type	N/A
Generic Specification Reference No.	N/A

Dungarvan Additional Information Panel				
Reel Unwind Direction	N/A			

text proof-read and approved. RSC GSK is responsible for site technical requirements and pre-press suitability.

is responsible to advise RSC in case changes required impact the followings:

Formulation Tablet embossing Storage conditions **Shelf Life**